

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number
WO 01/16367 A1

- (51) International Patent Classification⁶: C12Q 1/68 (74) Agent: BORTNER, Scott, R.; The Perkin-Elmer Corporation, 850 Lincoln Centre Drive, Foster City, CA 94404 (US).
- (21) International Application Number: PCT/US99/06978
- (22) International Filing Date: 26 March 1999 (26.03.1999) (81) Designated States (*national*): AU, CA, JP.
- (25) Filing Language: English (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data: 09/048,880 26 March 1998 (26.03.1998) US Published:
— With international search report.
- (71) Applicant: THE PERKIN-ELMER CORPORATION [US/US]; 850 Lincoln Centre Drive, Foster City, CA 94404 (US).
For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.
- (72) Inventors: AOYAGI, Kazuko; 3 Admiral Drive #457, Emeryville, CA 94608 (US). LIVAK, Kenneth, J.; 1148 Polk Lane, San Jose, CA 95117 (US).

(54) Title: METHODS FOR EXOGENOUS, INTERNAL CONTROLS DURING NUCLEIC ACID AMPLIFICATION



(57) Abstract: Reporter-quencher probe assays of nucleic acid amplification, such as PCR, are rendered more meaningful by the addition of internal control reagents. An internal control polynucleotide is amplified with internal control primers and the product is measured by correlation with increased fluorescence by polymerase mediated-exonuclease cleavage or hybridization of the internal control probe. Probes specific for target and internal control polynucleotides are labelled with spectrally resolvable reporters, allowing for concurrent detection and measurement of target and control amplification. A kit of all PCR reagents can be dispensed into reaction chambers in a high-throughput system for rapid and accurate nucleic acid amplification assay, with real-time or end-point measurements. Fluorescent signals correlated to target and internal control levels are spectrally resolvable and measured concurrently. A non-extending oligonucleotide or nucleic analog "block", complementary to the internal control polynucleotide, is added to the amplification mixture to preclude amplification of the internal control polynucleotide and function as an internal negative control. The amplification control reagents, kits, and methods of the present invention provide positive and negative control tests occurring within, and measurable within, the reaction chamber.

BEST AVAILABLE COPY

**METHODS FOR EXOGENOUS, INTERNAL CONTROLS
DURING NUCLEIC ACID AMPLIFICATION**

5

FIELD OF THE INVENTION

The invention relates generally to the field of nucleic acid amplification, and more particularly, to methods of internal controls that verify the absence or
10 presence of specific target sequences during the polymerase chain reaction.

REFERENCES

Agrawal, S. et al, "Site-specific functionalization of oligonucleotides for attaching
two different reporter groups", *Nucleic Acids Research*, 18: 5419-5423
15 (1990).

Ambion Technical Bulletin, Ambion Inc., <http://www.ambion.com>.

Andrus, A. in *Evaluating and Isolating Synthetic Oligonucleotides*, (1992),
Applied Biosystems, Inc., Foster City, CA.

Andrus, A., "Chemical methods for 5' non-isotopic labelling of PCR probes and
20 primers" (1995) in *PCR 2: A Practical Approach*, Oxford University Press,
Oxford, pp. 39-54

- Barany, F., "The Ligase Chain Reaction in a PCR World", PCR Methods and Applications 1:5-16 (1991).
- Barany, F. "Genetic disease detection and DNA amplification using cloned thermostable ligase", Proc. Nat. Acad. Sci USA 88:189-193 (1991).
- 5 Beaucage, S.L. and Iyer, R.P. "Advances in the synthesis of oligonucleotides by the phosphoramidite approach", Tetrahedron 48:2223-2311 (1992).
- Bergot et al, "Spectrally resolvable rhodamine dyes for nucleic acid sequence determination", US Patent 5,366,860, issued Nov. 22, 1994.
- Bevan et al, "Sequencing of PCR-amplified DNA", PCR Methods and
- 10 Applications, 1:222-228 (1992)
- Blackburn, G.M. and Gait, M.J. Eds. "DNA and RNA structure" in *Nucleic Acids in Chemistry and Biology*, 2nd Edition, (1996) Oxford University Press, pp. 15-81.
- Caruthers, M. and Beaucage, S., "Phosphoramidite compounds and processes"
- 15 US Patent No. 4,415,732, issued Nov. 15, 1983.
- Cardullo et al, "Detection of nucleic acid hybridization by non-radiative fluorescence resonance energy transfer", Proc. Natl. Acad. Sci., 85:8790-8794 (1988).
- Clegg, R., "Fluorescence resonance energy transfer and nucleic acids", Meth.
- 20 Enzymol., 211:353-388 (1992)
- Dieffenbach, C.W. and Dveksler, G.S. in *PCR Primer: A Laboratory Manual* (1995) Cold Spring Harbor Laboratory Press, pp. 13-14.

- Egholm, M, Buchardt, O. Christensen, L, Behrens, C, Freier, S., Driver, D.,
Berg, R., Kim, S. "PNA hybridizes to complementary oligonucleotides
obeying the Watson-Crick hydrogen bonding rules", *Nature*, 365:566-68
(1993).
- 5 Fung et al, "Method of detecting electrophoretically separated oligonucleotides",
U.S. patent 4,855,225, issued Aug. 8, 1989.
- Gelfand, D., Holland, P., Saiki, R., and Watson, R., "Homogeneous assay
system using the nuclease activity of a nucleic acid polymerase", US Patent
5,210,015, issued May 9, 1993.
- 10 Gilliland et al, "Analysis of cytokine mRNA and DNA: Detection and
quantitation by competitive polymerase chain reaction", *Proc. Natl. Acad.
Sci.*, 87:2725-2729 (1990)
- Green et al, "Sequenced-tagged site (STS) content mapping of human
chromosomes: Theoretical considerations and early experiences", *PCR
Methods and Applications*, 1:77-90 (1991).
- 15 Hermanson, G. T., in *Bioconjugate Techniques*, (1996) Academic Press, San
Diego, pp. 40-55, 643-671.
- Higuchi, R., Fockler, C., Dollinger, G. and Watson, R., "Kinetic PCR: Real time
monitoring of DNA amplification reactions" *Biotechnology*, 11:1026-30
20 (1993)
- Higuchi, R., Dollinger, G., Walsh, P., and Griffith, R., "Simultaneous
amplification and detection of specific DNA sequences" *Biotechnology*,
10:413-17 (1992)

- Holland, P.M., Abramson, R., Watson, R. and Gelfand, D., "Detection of specific polymerase chain reaction product by utilizing the 5' to 3' exonuclease activity of *Thermus aquaticus* DNA polymerase" *Proc. Natl. Acad. Sci.*, 88:7276-80 (1991).
- 5 Horn, T. and Urdea, M., "A chemical 5'-phosphorylation of oligodeoxyribonucleotides that can be monitored by trityl cation release", *Tetrahedron Lett.*, 27: 4705-08 (1986)
- Innis et al, in *PCR Protocols*, Academic Press, NY (1989)
- Ju, J. et al, "Design and Synthesis of fluorescence energy transfer dye-labeled
10 primers and their application for DNA sequencing and analysis", *Analytical Biochemistry*, 231:131-140 (1995).
- Keller, G. and Manak, M., in *DNA Probes, Second Edition*, (1993), Stockton Press, New York.
- Kricka, L., in *Nonisotopic DNA Probe Techniques* (1992), Academic Press, San
15 Diego, pp. 3-28.
- Lawyer, F.C., Stoffel, S., Saiki, R., Myambo, K., Drummond, R. and Gelfand, D., "Isolation, characterization, and expression in *Escherichia coli* of the DNA polymerase gene from the extreme thermophile, *Thermus aquaticus*." *J. Biol. Chem.* 264:6427-37 (1989).
- 20 Lee, L.G., Connell, C., and Bloch, W., "Allelic discrimination by nick-translation PCR with fluorogenic probes", *Nucl. Acids Res.*, 21:3761-66 (1993)
- Livak, K., Flood, S., Marmaro, J., Giusti, W., and Deetz, K., "Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system

- useful for detecting PCR product and nucleic acid hybridization". PCR Methods and Applications, 4:357-362 (1995)
- Livak, K. US Ser. 08/657,689 and WO 97/46708, "Passive internal references for the detection of nucleic acid amplification products"
- 5 Livak, K., Flood, S., Marmaro, J. "Method for Detecting Nucleic Acid Amplification Using Self-Quenching Fluorescence Probe", US Patent 5,538,848, issued July 23, 1996.
- Livak, K., Marmaro, J., and Todd, J., "Towards fully automated genome-wide polymorphism screening" Nature Genetics, 9:341-42 (1995).
- 10 Longo, M.C., Berninger, M.S., and Hartley, J.L. (1990) "Use of uracil DNA glycosylase to control carryover contamination in polymerase chain reactions" Gene, 93:125-28.
- Lyamichev, V., Brow, M., Dahlberg, J., "Structure-specific endonucleolytic cleavage of nucleic acids by eubacterial DNA polymerases". Science, 15 260:778-83 (1993).
- McPherson, M.J., Quirke, P., and Taylor, G.R. in *PCR 2: A Practical Approach* (1995) Oxford University Press, Oxford, pp. 7, 19.
- McPherson, M.J., Quirke, P., and Taylor, G.R. in *PCR: A Practical Approach, Volume 1* (1991) Oxford University Press, Oxford, pp. 46, 199.
- 20 Menchen et al, "4,7-Dichlorofluorescein dyes as molecular probes", U.S. patent 5,188,934, issued Feb. 23, 1993.

- Mullah B. and Andrus, A., "Automated synthesis of double dye-labeled oligonucleotides using tetramethylrhodamine (TAMRA) solid supports", *Tetrahedron Letters*, 38: 5751-5754 (1997).
- 5 Mullah, B., Livak, K., Andrus, A. and Kenney, P., "Efficient synthesis of double dye-labeled oligodeoxyribonucleotide probes and their application in a real time PCR assay", *Nucl. Acids Res.* 26:1026-1031 (1998).
- Nielsen, P.E., Egholm, M., Berg, R.H., and Buchardt, O., "Sequence-selective recognition of DNA by strand displacement with a thymidine-substituted polyamide", *Science* 254:1497-1500 (1991).
- 10 Orum, H., Nielsen, P., Egholm, M., Berg, R., Buchardt, O., and Stanley, C. "Single base pair mutation analysis by PNA directed PCR clamping", *Nuc. Acids Res.* 21:5332-36 (1993).
- Ozaki, H. and McLaughlin, L., "The estimation of distances between specific backbone-labeled sites in DNA using fluorescence resonanace energy transfer", *Nucleic Acids Research*, 20: 5205-5214 (1992)
- 15 Perseptive Biosystems, "Identifying point mutations by PNA-directed PCR clamping" (1995).
- Ruth, J., "Single-stranded labelled oligonucleotides, reactive monomers and methods of synthesis" U.S. patent 4,948,882, issued Aug. 14, 1990.
- 20 Theisen, P., McCollum, C., and Andrus, A. "Fluorescent dye phosphoramidite labelling of oligonucleotides", *Nucleic Acid Symposium Series No. 27*, Oxford University Press, Oxford, pp. 99-100 (1992).

Tsai S-J, and Wiltbank M.C., "Quantification of mRNA Using Competitive RT-PCR with Standard Curve Methodology" *Biotechniques* 21:5:862-866 (1996).

5 Tyagi, S. and Kramer, F.R., "Molecular Beacons: Probes that fluoresce upon hybridization", *Nature BioTechnology*, 14:303-08 (1996).

Tyagi, S. and Kramer, F.R., "Detection probes, kits, and assays" *Intl. Publ. WO* 97/39008.

Urdea et al, "Nucleic acid probes" U.S. patent 5,093,232, issued Mar. 3, 1992.

10 Van der Laan, A. et al, "A Convenient automated solid-phase synthesis of PNA-(5')-DNA-(3')-PNA chimera", *Tetrahedron Lett.*, 38:2249-52 (1997).

Vinayak, R. et al, "Automated chemical synthesis of PNA-DNA chimera on a nucleic synthesizer", *Nucleosides & Nucleotides*, 16:1653-56 (1997).

15 Walker, G. et al, "Strand displacement amplification - an isothermal, *in vitro* DNA amplification technique", *Nucleic Acids Research*, 20: 1691-1696 (1992).

Walker, G., Little, M., Nadeau, J. and Shank, D., "Isothermal *in vitro* amplification of DNA by a restriction enzyme/DNA polymerase system" *Proc. Natl. Acad. Sci.* 89:392-96 (1992).

20 Wang, M., Doyle, M.V., and Mark, D.F., "Quantitation of mRNA by the Polymerase Chain Reaction", *Proc. Natl. Acad. Sci. USA* 86:9717-9721 (1989).

Woo et al., "Rhodamine phosphoramidite compounds", U.S. Patent No. 5,231,191, issued July 27, 1993.

Woo, S. and Fung, S., "Solid support reagents for the synthesis of 3'-nitrogen containing polynucleotides" US Patent No. 5,552,471, issued Sept. 3, 1996.

Zimmermann, K., Schogl, D., Plaimauer, B., and Mannhalter, J.W., Quantitative
5 Multiple Competitive PCR of HIV-1 DNA in a Single Reaction Tube",
BioTechniques 21:481-484 (1996).

BACKGROUND

Nucleic acid amplification, and the polymerase chain reaction (PCR) in
10 particular, has become an important research tool, with applications in cloning,
analysis of genetic expression, DNA sequencing, genetic mapping, drug
discovery, and the like (Gilliland et al, 1990; Bevan et al, 1992; Green et al, 1991).
Descriptions of, and guidance for conducting, PCR is provided in extensive
literature on the subject (Innis et al, 1989; McPherson et al, 1991; McPherson
15 et al, 1995).

Automated instrumentation has been developed for carrying out nucleic
acid amplifications, most commonly automated thermal cyclers for conducting
PCR and the like. Important design goals fundamental to PCR instrument
development have included fine temperature control, minimization of sample-to-
20 sample variability in high-throughput, highly multiplexed-sample thermal cycling,
automation of pre- and post-PCR processing steps, high-speed cycling,
minimization of sample volumes, real time measurement of amplification
products, minimization of cross-contamination or sample carryover, and the like.

In particular, the design of instruments that permit PCR to be carried out and monitored in real time is highly desirable. Reaction chambers that remain closed during amplification and analysis are desirable for preventing cross-contamination (Higuchi et al, 1992; Higuchi et al, 1993; Holland et al, 1991).

5 Methods, reagents, and kits of reagents that facilitate multi-channel pipetting or robotic dispensing are desirable. A limited number of automated pipetting and dispensing steps will minimize errors and ambiguous results. Clearly, the successful realization of these design goals would be especially desirable in the analysis of diagnostic samples, where a high frequency of false
10 positives and false negatives would severely reduce the value of the PCR-based procedure.

 Fluorescence-based approaches to provide real time measurements of amplification products during a PCR (Holland et al, 1991) have either employed intercalating dyes (such as ethidium bromide) to indicate the
15 amount of double stranded DNA present (Higuchi, 1992; Higuchi, 1993; Gelfand et al, 1993) or probes containing reporter-quencher pairs ("TaqMan®", exonuclease assay) that are cleaved during amplification to release a fluorescent signal that is proportional to the amount of double stranded DNA present (Livak 1996). The polymerase that conducts primer
20 extension and amplifies the polynucleotide also possesses a 5'→3' exonuclease activity that serves to cleave the probe. In the exonuclease assay, a "reporter" dye and a "quencher" dye are attached to an oligonucleotide probe which is complementary to the target DNA. The dyes

are selected and arranged to interact through a fluorescence resonance energy transfer (FRET) process (Clegg, R., 1992). The reporter is a luminescent compound that can be excited either by chemical reaction, producing chemiluminescence, or by light absorption, producing fluorescence (Figure 1). The quencher can interact with the reporter to alter its light emission, usually resulting in the decreased emission efficiency of the reporter. This phenomenon is called quenching. The efficiency of quenching is a strong function of the distance between the reporter molecule and the quencher molecule. Thus, in a nucleic acid hybridization assay, detection of a hybridization event is accomplished by designing an energy transfer system in which the spacing between a reporter and a quencher is modulated as a result of the hybridization. Two examples of systems that perform the exonuclease assay and other quantitation, fluorescent-based arrays are the ABI PRISM™ 7700 and ABI PRISM™ 7200 Sequence Detection Systems (Perkin-Elmer).

The exonuclease assay of nucleic acid amplification employing reporter-quencher probes (Lee et al, 1993; Livak et al, 1995) gives direct detection of polymerase chain reaction (PCR) products with no downstream sample processing. The quencher is released from its close proximity to the reporter upon cleavage so that the signal from the reporter is no longer quenched. An increase in fluorescence occurs which correlates directly and proportionally with the increase in copies of the PCR product. By using real-time or end-point analysis, detection and quantitation of PCR products can be obtained by

measuring the increase in fluorescence of cleaved, self-quenching fluorescent probes.

Release of fluorescence can be detected and measured by laser-induced fluorescence with an optical-fiber probe in a non-invasive, closed reaction chamber. The assay is conducted by a high-throughput, data sampling routine during the course of PCR thermal cycling (real-time analysis) or at the end of PCR (end-point analysis). Automated quantitation using PCR is highly desirable. Manual methods rely on end-point electrophoresis of an aliquot of the PCR and spectroscopic or densitometric quantitation. Real time monitoring of PCR permits far more accurate quantitation of starting target DNA concentrations in multiple-target amplifications than manual end-point methods. With calibrated, internal controls, the relative values of close concentrations can be resolved by factoring the history of the relative concentration values during the PCR. Real time monitoring of target with internal controls is desirable.

Internal and/or parallel tests that confirm conditions for amplification of the target are desirable as control tests. When used in a high-throughput format, such as 96 well, microtitre plate configurations, PCR often is plagued by false positives due to template contamination from adjacent wells, pipetting errors, or aerosol transmission. In addition, PCR suffers from false negative results when enzyme inhibitors are present in the target samples or when reagents are missing or degraded. Therefore, control amplification tests are desirable.

Positive amplification control tests give a detectable product derived from a component that is separate and distinct from the target. Detection of the

positive control product indicates that amplification is viable and operative within the reaction chamber. Positive amplification control tests which give no detectable product from the control components, indicate conditions within the reaction chamber that do not allow amplification. Another desirable feature of amplification control is to "turn off" the positive control signal with the addition of a negative or "blocking" element to other reaction chambers during the assay, allowing the measurement of background.

Internal amplification controls are to be distinguished from "passive" internal reference molecules (Livak, Ser. 08/657,689) which provide for signal and detection corrections. Passive internal references, such as non-complementary, reporter-quencher molecules do not hybridize to target or other polynucleotides, are not consumed or act as substrates for enzymes, and do not undergo chemical or enzymatic reactions of any sort. Thus, passive internal references do not provide verification or indication of conditions for amplification, within the reaction chamber.

In addition to the above limitations and problems associated with signal and detection, a passive internal reference does not address the very common issues of, (i) contamination of target or other reagents with foreign DNA, (ii) inhibition of PCR, and (iii) confirming amplification efficiency within the reaction chamber. Reporter-quencher probe assays with internal fluorescence-generating controls are needed to provide accurate, precise, and sensitive measurements of changes in fluorescence that are attributable solely to formation of the amplification product.

Control amplification reactions are necessary for (i) normalization of quantitation results, (ii) detection of amplification inhibitors in the target and other reagents, and (iii) to establish background signal levels. A pervasive difficulty is keeping amplification of the control polynucleotide from interfering with target amplification or detection of the product. Internal control polynucleotides (ICP) undergo amplification within the same reaction chamber as the known or unknown target polynucleotide, imparting convenience in preparing samples and measuring results. The ICP may be endogenous, i.e. from the same source, genome, chromosome, gene, plasmid, or fragment as the target.

10 Endogenous ICP are subject to amplification inhibitors and can therefore give a false negative signal. Endogenous ICP also may have priming sites for target primers and therefore give a false positive signal. In fact, endogenous ICP systems may share one or more primers with the target. Exhaustion of shared primers leads to inaccurate PCR quantitation and limited dynamic range.

15 Another negative feature of endogenous ICP is the necessity to select, design, and purify ICP, ICP primers, and ICP self-quenching probe for each target to ensure compatibility and viable amplification. A universal exogenous ICP which is not derived from the same source, etc. as the target is therefore desirable, to avoid these disadvantages.

20 In addition to the positive control amplification, it is desirable to have separate, parallel negative control tests to establish a background signal level for quantitative systems such as the exonuclease assay. By "switching off" the control amplification, the fluorescence of the intact self-quenching probe of the

internal control polynucleotide can be monitored and subtracted as a baseline value from samples undergoing normal amplification. Previous attempts at negative control background measurement have entailed; (i) deletion of essential components of amplification, such as magnesium, dNTP, and enzyme, or (ii) 5 addition of amplification inhibitors such as sodium dodecyl sulfate (SDS) or EDTA. All of these attempts suffer from the disadvantage of changing, distorting, or obscuring fluorescent signal generation or detection.

Most techniques used for total RNA isolation yield RNA with significant amounts of genomic DNA contamination (Ambion). Reverse transcription, 10 polymerase chain reaction (RT-PCR) is a popular method for analyzing low abundance mRNA from limiting amounts of tissue and the quantitative analysis of gene expression. Because the PCR step involves an exponential amplification, small tube-to-tube variations in amplification efficiency can translate into dramatic differences in the yield of final product and gross errors in 15 estimation of initial abundance. A frequent cause of concern among investigators performing quantitative RT-PCR is inaccurate data caused by DNA contamination in RNA preparations. PCR cannot discriminate between cDNA targets synthesized by reverse transcription and genomic DNA contamination. Although DNA contamination is easily detected by performing a 'no-RT' 20 negative control, there is no satisfactory and comprehensive solution to contamination and inhibition which the present invention provides. The conventional methods of RT-PCR quantitative analysis (Wang et al, 1989; Tsai et al, 1996; Zimmermann et al, 1996) are all gel-based assays, which rely on

imprecise and subjective visualization techniques.

In view of the limitations and deficiencies of conventional controls for the quantitation and detection of nucleic acid amplification products, it is of interest to develop non-gel based internal control methods that provide both negative
5 and positive indications amplification. The invention described herein provides for such internal control reagents, kits, and methods.

SUMMARY

The present invention is directed towards novel methods and a kit of reagents for quantitating nucleic acid amplification of control DNA in the
10 presence of, and concurrently with, nucleic acid amplifications of known and unknown target DNA.

It is an object of certain embodiments of the present invention to provide a method of conducting nucleic acid amplification reactions in a single reaction chamber whereby internal control primers hybridize to an internal control
15 polynucleotide and target primers hybridize to a target polynucleotide, and amplification of polynucleotides occurs by polymerase chain extension, thermal melting, and hybridization. Internal control polynucleotides are comprised of DNA or RNA and are not sequence-homologous to target polynucleotides. Internal control polynucleotides do not allow amplification by target primers or
20 hybridization with target probes. The amplification product resulting from amplification of the internal control polynucleotide with internal control primers will typically be from 50-500 bp in length. The amplicon will contain an internal

site complementary for hybridization of the internal control probe. An internal control polynucleotide comprised of RNA may be a substrate for reverse-transcriptase and result in the production of a cDNA copy.

It is another object of certain embodiments of the present invention to provide
5 internal control reagents and methods that detect the presence of PCR inhibitors in the target and other components in the PCR mixture. The internal controls will have the properties of being easy to dispense, i.e. a minimum of pipetting operations and conducive to robotic automation in the conventional 96-well microtitre plate format. The internal control reagents should not decrease PCR amplification efficiency of the
10 target polynucleotide. Both negative and positive internal control signals should give distinct and operative results. A positive internal control signal can discriminate a true negative target signal from a false negative. Measurement of a negative internal control signal gives the background which is subtracted from the positive internal control signal. The fluorescence of the reporter on the internal control
15 probe is measured so as to normalize the reaction for such amplification factors that vary from reaction to reaction. Thus by examining the fluorescent signal from the internal control probe, the effects of most sources of PCR inhibition are identified.

Reporter-quencher probe assays including the subject internal control reagents may be used in conjunction with a variety of nucleic acid amplification systems.
20 Generally, the assays require either the use of a nucleic acid polymerase with exonuclease activity or a population of double stranded DNA which increases during the course of the reaction being monitored. Exemplary amplification schemes that may be employed with the system of the invention include PCR, ligase-based

amplification schemes, such as ligase chain reaction, LCR (Barany, 1991), Q-beta replicase-based amplification schemes, strand displacement amplification (SDA) schemes (Walker et al, 1992; Keller, 1993).

An exogenous ICP is selected and designed to have no sequence
5 homology with any known target from inspection of sequence databanks. The exogenous ICP is DNA or RNA of 50-500 bp in length. The exogenous ICP is added to the amplification reagent kit, along with ICP primers and self-quenching probe, in known amounts to consistently yield a known amount of fluorescent signal. The exogenous ICP can verify amplification conditions
10 within a reaction chamber to establish a true negative for the presence of target DNA.

It is yet another object of certain embodiments of the present invention to provide self-quenching fluorescence probes with different and non-homologous sequences for hybridization to the internal portions of target and internal control
15 polynucleotides. Since the internal control polynucleotide (ICP) is designed or selected to have minimal homology with any known target polynucleotide, the internal control probe will hybridize only to ICP and not to target polynucleotide. Similarly, the target probe will hybridize only to target polynucleotide, and not to ICP. Self-quenching probes exist in at least one
20 single-stranded conformation when unhybridized such that the quencher dye quenches the fluorescence of the reporter dye. When hybridized to polynucleotides, the probes exist in at least one conformation where the fluorescence of the reporter dye is unquenched and the fluorescence intensity of


reporter dye is greater than the fluorescence intensity of the quencher dye when the probe is hybridized to polynucleotide. Nucleic acid polymerase can substantially digest the probes during amplification to separate the reporter dye from the quencher dye. Fluorescence from the cleaved reporter dye corresponds to the occurrence of nucleic acid amplification. Fluorescence may also accrue based on hybridization of the probe to complementary target or ICP. The target and internal control probes are labelled with different and spectrally resolvable reporter moieties when used in the same reaction chamber. The respective emission spectra from the reporter moieties with a reaction chamber must be sufficiently non-overlapping so that separate emission contributions can be resolved. The separate peaks may be quantitated, correlating to the relative amounts of amplification products, i.e., target and internal control polynucleotides.

Clearly, the system may be generalized to include a plurality of fluorescent reporters, e.g. to monitor the simultaneous amplification of several target nucleic acids in a single reaction, so that a plurality of fluorescent intensity ratios are monitored. Such multi-reporter systems are advantageous in applications requiring the analysis of multiple amplifications occurring in a single reaction chamber. In such systems, each of the reporter molecules produce emissions which are spectrally resolvable from the emissions from any of the other reporters. The particular quencher used with each reporter can be the same or different, depending on the spectral properties of the quencher and reporter. Several spectrally resolvable dyes

suitable for use in such embodiments are disclosed in Fung et al; Menchen et al; Bergot et al; and like references.


In a preferred embodiment of the self-quenching fluorescence probe, the reporter dye is separated from the quencher dye by at least 12 nucleotides, the
5 reporter dye is attached at the 5' terminus or 3' terminus of the self-quenching fluorescence probe, and the quencher dye is attached at the 5' terminus or 3' terminus. In another preferred embodiment, a first self-quenching fluorescence probe is complementary to the internal control polynucleotide and a second self-quenching fluorescence probe is complementary to the target polynucleotide.
10 Additionally, the reporter dye of the first self-quenching fluorescence probe is spectrally resolvable from the reporter dye of the second self-quenching fluorescence probe. Preferably, in this embodiment, an excitation beam is generated from the 488 nm emission line of an argon ion laser to induce fluorescence in the reporter dye. Fluorescence emission from the reporters and
15 quenchers can be detected from 500 to 650 nm. Fluorophores in a reaction chamber can be discriminated when the emission maxima are sufficiently different and the emission bandwidth is sufficiently narrow.

Preferred embodiments of reporter moieties are fluorescein dyes with the general structure and numbering system below, where L is a linker.



Preferred embodiments of fluorescein reporter dyes are selected from the group
 5 consisting of 5-carboxyfluorescein (5-FAM), 6-carboxyfluorescein (6-FAM),
 2',4',1,4,-tetrachlorofluorescein (TET), 2',4',5',7',1,4-hexachlorofluorescein
 (HEX), and 2',7'-dimethoxy-4',5'-dichloro-6-carboxyrhodamine (JOE)
 (Figure 2). Other embodiments of reporter moieties are cyanine dyes, dansyl
 derivatives, and the like.

Preferred embodiments of quencher moieties are; (i) rhodamine dyes selected from the group consisting of tetramethyl-6-carboxyrhodamine (TAMRA), and tetrapropano-6-carboxyrhodamine (ROX)., and (ii) DABSYL, DABCYL, cyanine, anthroquinone, nitrothiazole, and nitroimidazole compounds and the like (Figure 3). Rhodamine dyes may bear the general structure and numbering system below, where L is a linker.



Fluorescein and rhodamine derivatives of the present invention may be substituted at one or more of the numbered positions above.

It is another aspect of the present invention to provide an oligonucleotide or nucleic acid analog, referred hereafter as a "block", that is complementary to the ICP and which is not extendable during nucleic acid amplification. The block may hybridize to any portion of the ICP with an equal or greater affinity than the internal control primers and prevent amplification of the ICP. Addition of the block to PCR will effect a negative result to the internal control. No amplification product from the ICP will result, as detected by the absence or minimal background of fluorescence from the internal control probe. Fluorescence from the internal control probe is measured at the emission

maxima, e.g., occurring between 500-650 nm of the internal control reporter. The reporter on the target probe is chosen to have little or no spectral overlap with the reporter of the internal control probe. Therefore the reporter on the target probe can be measured independently from and concurrently with the reporter on the internal control probe within a single reaction chamber.

The block may be comprised of modifications to the internucleotide linkage, the sugar, or nucleobase moieties of a DNA primer to render it non-extendable by polymerase. An example of a suitable modification is a 3' phosphate group. Analogs of DNA may be employed as the block, such as, 2-aminoethylglycine, peptide-nucleic acid (PNA) and other amide-linked oligomers; 2'-O-methyl and other 2'-O-alkyl oligoribonucleotides; phosphorothioate and other phosphate analogs; and the like. The block is selected for several properties, including (i) high specificity, (ii) high affinity, (iii) non-extendability, (iv) chemical stability, (v) non-interference with amplification. In a preferred embodiment, the block is a PNA (peptide-nucleic acid) oligomer (Nielsen, P.E. et al). To improve solubility and lower aggregation effects, the PNA block may be conjugated with hydrophilic labels, such as polyethyleneoxy, peptides, nucleic acids, nucleic acid analogs, and the like.

PNA has been used in PCR as a "clamping element" for single base-pair mutation analysis (Orum et al). In this report, PNA suppresses amplification of complementary target sequences, typically wild-type, and allows the selective amplification of low-copy number or mutant target sequences with competing DNA primers.

Certain preferred embodiments of the present invention include methods for the end-point and real-time measurements of amplification product formation. In an end-point mode, the fluorescence measurement is performed after the amplification reaction is complete, e.g., after all or substantially all of the amplification cycles of a PCR reaction have been completed. In a real-time mode, fluorescence measurements is performed multiple times during the amplification reaction, e.g., after each thermocycle of a PCR process. The real-time mode is preferred when a quantitative measure of the initial amount of target nucleic acid is required, e.g., the copy-number of pathogen nucleic acid present in a blood sample.

Fluorescent measurements may be used to verify the operability of PCR in the target vessel. The use of the internal control reagents of the invention permits the assignment of a near-zero background level of fluorescence for high-sensitivity tests, e.g. low copy-number or rare genes. The use of the internal control reagents of the invention permits the simultaneous use of multiple reporter-quencher probes in reporter-quencher probe assays.

It is an object of the invention to provide a method for accurate, real time monitoring of nucleic amplification reactions by providing apparatus and fluorescent reagents for generating a stable fluorescent signal proportional to the amount of amplification product. The availability of data showing the progress of amplification reactions leads to more accurate estimates of relative starting concentrations of target nucleic acids, to rapid assessment of the efficiency of the amplification reactions, and opens the possibility of reduced reagent usage and

feedback reaction control.

It is an object of the present invention to provide a kit consisting of reagents required for nucleic acid amplification reactions for practicing the improved amplification methods of the invention. Kits make the practice of the claimed methods more reproducible and easier to perform. Kits may supply
5 reagents in pre-measured amounts so as to simplify the performance of the subject methods. Furthermore, kits typically contain detailed instructions for carrying out the methods of the invention. In one embodiment of the invention, the kit comprises an internal control polynucleotide, internal control primers, a
10 non-extending oligonucleotide or nucleic acid analog complementary to the internal control polynucleotide, target primers, a nucleic acid polymerase having 5'→3' nuclease activity, self-quenching fluorescence probes including reporter and quencher moieties, 5' nucleotide triphosphates. The kits of the invention may further comprise additional reagents that are necessary for performing the
15 subject methods, such reagents include, but are not limited to uracil N-glycosylase (UNG), buffers, molecular size standards, wax beads, and the like.

It is an object to employ the methods of the invention to sample tracking and signal calibration at the point of target sample extraction, isolation, or purification. For these purposes, internal control reagents comprised of ICP,
20 internal control primers and probe may be added to crude sample preparations.

It is an object to employ the methods of the invention to reverse-transcriptase/PCR on mRNA samples, especially for low copy-number genes.

It is an object to employ the methods of the invention to allelic discrimination of polymorphic samples. Variations of a gene can be distinguished at the level of single base-pair differences of inherited alleles by the exonuclease assay (Lee et al, 1993).

5 It is an object to employ the methods of the invention to plus/minus, target-specific assays for pathogen detection, whereby one pair, or a limited plurality of pairs, of primers and a single reporter-quencher target probe are components of an amplification kit. The kit will also contain the internal control reagents and other components necessary for amplification. Measured aliquots
10 from the kit can be dispensed in a high-throughput or automated manner to locations in a spatially-addressable test holder. Locations can be wells, sites, or surface arrays. The preferred embodiment of the holder is a microtitre well tray. The preferred embodiment of the locations are wells in a microtitre well tray. Typical location density is 96 wells in an 8x12 configuration of wells conforming
15 to the accepted industry standards. Materials for the locations of the holder may be polymeric, metallic, glass, or ceramic. Target samples can be dispensed in a high-throughput or automated manner to the locations before or after delivery of the kit components.

BRIEF DESCRIPTION OF THE DRAWINGS

- 20 FIG. 1 Exonuclease assay target polynucleotide with self-quenching probe
FIG. 2 Fluorescein reporter structures: FAM, TET, HEX, JOE
FIG. 3 Quencher structures: TAMRA, ROX, DABCYL, DABSYL

- FIG. 4 Exonuclease assay of internal control polynucleotide; positive and negative control aspects
- FIG. 5 Chart of reporter and quencher spectral properties: emission and absorbance maxima
- 5 FIG. 6 Self-quenching probe structure: 5' FAM reporter and 3' TAMRA quencher
- FIG. 7 Exonuclease assay results and sample assignment in 96-well format, with internal negative (A1-A6) and positive (A7-A12) control wells.
- FIG. 8 Detection of E. Coli 0157:H7. Exonuclease assay results from
- 10 Example 4.
- FIG. 9 Detection of Mycoplasma synoviae DNA. Exonuclease assay results from Example 5
- o Well: location in 96 well format (Figure 7)
 - o Type: NAC - no amplification control, NTC - no template control,
 - 15 Unknown
 - o Sample: target copy number by dilution, or negative
 - o PCR: positive (+) or negative (-) result
 - o Rn: reporter minus NTC (A7-A12)
 - o Ct: threshold cycle number; PCR cycle number at which fluorescence
 - 20 is distinguishable from background

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Reference will now be made in detail to the preferred embodiments of the invention, examples of which are illustrated in the accompanying drawings. While the invention will be described in conjunction with the preferred embodiments, it will be understood that they are not intended to limit the invention to those embodiments. On the contrary, the invention is intended to cover alternatives, modifications, and equivalents, which may be included within the invention as defined by the appended claims.

10 I. DEFINITIONS

Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings:

“Polynucleotide” or “oligonucleotide” refer to linear polymers of natural nucleotide monomers or analogs thereof, including double and single stranded deoxyribonucleotides “DNA”, ribonucleotides “RNA”, α -anomeric forms thereof, and the like. In other words, an “oligonucleotide” is a chain of deoxyribonucleotides or ribonucleotides, that are the structural units that comprise deoxyribonucleic acid (DNA) and ribonucleic acid (RNA), respectively.

20 “Nucleotide” is the monomer unit in biopolymer nucleic acids, such as DNA or RNA. A nucleotide is composed of three moieties: sugar, phosphate, and nucleobase (Blackburn, M., 1996). When part of a duplex, nucleotides are

also referred to as "bases" or "base pairs". The most common naturally-occurring nucleobases, adenine (A), guanine (G), uracil (U), cytosine (C, and thymine (T) bear the hydrogen-bonding functionality that binds one nucleic acid strand to another in a sequence specific manner. "Nucleoside" refers to a nucleotide that lacks a phosphate dye. Usually the nucleoside monomers are linked by phosphodiester linkages, where as used herein, the term "phosphodiester linkage" refers to phosphodiester bonds or bonds including phosphate analogs thereof, including associated counter-ions, e.g., H^+ , NH_4^+ , Na^+ , and the like. Polynucleotides typically range in size from a few monomeric units, e.g. 8-40, to several thousand monomeric units. Most molecular biology applications for polynucleotides require unique sequences of 15-30 nucleotides in length. Whenever a DNA polynucleotide is represented by a sequence of letters, such as "ATGCCTG," it will be understood that the nucleotides are in 5'→3' order from left to right and that "A" denotes deoxyadenosine, "C" denotes deoxycytidine, "G" denotes deoxyguanosine, and "T" denotes thymidine, unless otherwise noted.

"Watson/Crick base-pairing" and "Watson/Crick complementarity" refer to the pattern of specific pairs of nucleotides, and analogs thereof, that bind together through hydrogen-bonds, e.g. A pairs with T and U, and G pairs with C.

"Oligonucleotide analogs" are polymeric analogs of DNA and RNA made by chemical synthesis from monomeric nucleotide analog units, and possessing some of the qualities and properties associated with nucleic acids.

"Attachment site" refers to the atom on an oligonucleotide to which is attached the linker.

"Linker" refers to one or more atoms comprising a chain connecting an oligonucleotide and a label.

5 "Chimera" as used herein refers to an oligonucleotide including one or more nucleotide and one or more nucleotide analog units. The monomer units are linked through phosphodiester and phosphodiester analog linkages.

10 "Phosphodiester analog" refers to analogs of natural phosphodiester 3',5'-internucleotide linkages differing in their composition and/or location of attachment to a nucleotide, including but not limited to 2',5'-linkage, 3',3'-linkage, 5',5'-linkage, methyl phosphonate, alkylated phosphotriester, 3'-N-phosphoramidate, and PNA.

"Lower alkyl", "lower alkylene" and "lower substituted alkylene" refers to straight-chain, branched, or cyclic groups consisting of 1-12 carbon atoms.

15 "Label" refers to a group covalently attached at one or both termini of the nucleobase oligomer. The label is capable of conducting a function such as giving a signal for detection of the molecule by such means as fluorescence, chemiluminescence, and electrochemical luminescence (Kricka, L.). Alternatively, the label allows for separation or immobilization of the molecule
20 by a specific or non-specific capture method (Andrus, A., 1995).

"Energy transfer" and "fluorescence quenching" refer to a processes whereby energy is removed from an electronically excited luminescent "reporter" molecule by a "quencher" molecule, thereby returning the

reporter molecule to its ground state without the emission of light from the reporter molecule. The reporter molecule may be excited to one of its higher energy levels by any of a number of process, including light absorption and chemical reaction.

5 "Spectral resolution" and "spectrally resolvable" in reference to a plurality of dyes means that the fluorescent emissions of the dyes are sufficiently distinct, i.e., sufficiently non-overlapping, that reagents to which the respective dyes are attached, e.g. polynucleotides, can be resolved on the basis of a fluorescent signal generated by the respective dyes using standard
10 photodetection systems.

"Detection" refers to detecting, observing, or measuring a nucleobase oligomer on the basis of the properties of a covalently-attached detection label. Detection labels include, but are not limited to, fluorescent dyes, such as fluorescein and rhodamine derivatives, cyanine dyes, and energy-transfer dyes
15 (Clegg, R., 1992; Cardullo, 1988).

"Primer" refers to an oligonucleotide capable of selectively annealing to a specified target nucleic acid and thereafter serving as a point of initiation of a primer extension reaction wherein the primer is extended in a 5' → 3' direction.

20 "Primer extension reaction" refers to a reaction between a target/primer duplex and a nucleotide which results in the addition of the nucleotide to a 3'-end of the primer such that the added nucleotide is complementary to the corresponding nucleotide of the target nucleic acid.

The term "5'→3' nuclease activity" refers to an enzyme activity that cleaves nucleic acid at phosphodiester bonds. This activity can be either endo (cleaves at internal phosphodiester bonds) or exo (cleaves at the phosphodiester bond closest to either the 5' or 3' terminus of the nucleic acid strand).

- 5 The term "block" refers to a non-extendable oligonucleotide or nucleic acid analog complementary to the internal control polynucleotide. A block does not conduct polymerization or extension in the normal manner through extension of the 3' terminus of a DNA primer, with polymerase and deoxynucleotide triphosphates. A block functions as a negative control element, essentially
- 10 turning off amplification of the internal control polynucleotide and thereby precluding an increase in fluorescence derived from the internal control probe.

The term "self-quenching" refers to an intermolecular, energy transfer effect, e.g. a reporter and quencher are joined on an oligonucleotide in a configuration that permits energy transfer from the fluorophore to the quencher.

- 15 The term "end-point analysis" refers to a test where data collection occurs only when a reaction is complete. End-point analysis of the exonuclease assay entails reporter signal measurement when PCR is complete. Results are reported in terms of the change in fluorescence of the reporter signal, minus the internal control amplification fluorescence change.

- 20 The term "real-time analysis" refers to periodic monitoring during a test. Real-time analysis of the exonuclease assay measures reporter signal changes from cycle-to-cycle, minus the internal control amplification fluorescence change.

II. EXONUCLEASE ASSAY WITH INTERNAL CONTROL POLYNUCLEOTIDE AND BLOCK

The TaqMan® exonuclease assay (Holland et al, 1991; Lee et al; Livak, 1996) illustrated in Figure 1 shows self-quenching probe 1, including both a reporter label, F₁, and a quencher label, Q, and target primers 3a and 3b hybridized to target polynucleotide 2. During the polymerization phase of amplification, the primers 3a and 3b are then extended using a polymerase enzyme thereby forming extended primers 4a and 4b, e.g., using a DNA polymerase. During the primer extension reaction, a 5'→3' nuclease activity of the polymerase serves to cut the probe 1 so as to form probe fragments, including reporter-bearing fragment 5 and quencher-bearing fragment 6. Thus, the reporter and quencher labels are separated thereby preventing energy transfer between the two and the emission of the reporter becomes unquenched upon digestion of the probe. An increase in fluorescence, detectable as F₁, results.

When the internal control reagents and method of the present invention are practiced during the exonuclease assay, reagents additional to those above are added to the reaction chamber, including self-quenching probe 7 and primers 8a and 8b complementary to internal control polynucleotide 9 (Figure 4). During the polymerization phase of amplification, the primers 8a and 8b are then extended using a polymerase enzyme thereby forming extended primers using a

DNA polymerase. During the primer extension reaction, a 5'→3' nuclease activity of the polymerase serves to cut the probe 7 so as to form probe fragments, including reporter-bearing fragment 10 and quencher-bearing fragment 11. Thus, the reporter and quencher labels are separated thereby preventing energy transfer between the two and the emission of the reporter becomes unquenched upon digestion of the probe. An increase in fluorescence, detectable as F₂, results. Reporter dyes F₁ and F₂ are chosen to be spectrally resolvable. Quencher moieties on 1 and 7 may be the same or different.

When the negative internal control block 12 is added, no amplification of internal control polynucleotide 9 occurs. Non-extendable block 12 selectively binds to the 8a primer binding site, or at another site on 9, and precludes amplification. The self-quenching probe 7 remains intact and quenched. A negative control baseline may be measured and applied to positive control fluorescence changes.

15 III. DESIGN AND SYNTHESIS OF REAGENTS

Generally, the design and synthesis of oligonucleotides of the invention follows conventional teachings. Preferably, oligonucleotides are synthesized on an automated, solid-phase DNA synthesizer using phosphoramidite chemistry (Beaucage et al, 1992; Caruthers, 1983), e.g. Model 392 or 394 DNA synthesizer (PE Applied Biosystems, Foster City, CA).

a. Self-quenching probes

In designing self-quenching probes, the following general guidelines can be followed: (1) if the target nucleic acid sequence is located within a PCR amplicon, the probe sequence should be such that the probe hybridizes at a location on the sequence between the PCR primers; (2) probes should be about 20-30 nucleotides long so as to ensure good hybridization kinetics and specificity of binding; (3) avoid secondary structure in the probe and target nucleic acid sequence; (4) the probe should not hybridize to either of the forward and reverse primers; and (5) avoid probes with long stretches of a single nucleotide, i.e., more than four; and (6) when choosing between a probe sequence and its complement, pick the strand that has more C nucleotides than G nucleotides.

The self-quenching probe is designed so as to bring the reporter into close proximity with the quencher so as to permit efficient energy transfer from the reporter to the quencher. Guidance concerning the selection of an appropriate distance for a given embodiment is found in numerous references on resonant energy transfer between fluorescent reporter molecules and quenching molecules (also sometimes referred to as "donor" molecules and "acceptor" molecules, respectively), e.g. Clegg, 1992; Cardullo, 1988; Ozaki et al; Livak et al, 1995, and the like. Self-quenching probes may have intramolecular Watson/Crick base-pairing properties, i.e., self-complementarity. Adoption of a stable hydrogen-bonded conformation may result in a high-degree of energy transfer between the reporter and the quencher due to their enforced proximity (Tyagi et al, 1996; Tyagi et al, 1997). The increase in fluorescence of self-complementary,

self-quenching probes ("Molecular Beacons") upon hybridization to target or internal control polynucleotides during amplification may be significant enough to not require exonuclease cleavage of the probes.

The reporter and quencher are preferably close enough so that substantially all, e.g. 90%, of the fluorescence from the reporter is quenched. Typically, for energy transfer-based quenching, the distance between the first and second fluorophores should be within the range of 10-100 angstroms. Preferably, the fluorescer and quencher are separated by between about 4 to 10 nucleotides. However, the invention includes embodiments in which the number of nucleotides separating the fluorophores may be greater than 10. Preferably, either the reporter and quencher are attached to the 5' terminal nucleotide of the oligonucleotide probe. The reporter and quencher may also be attached to the 3' terminal nucleotide. In other embodiments of the reference molecules of the invention, the fluorescer and quencher are attached at internal sites on the polynucleotide. The invention also includes embodiments in which one of the two fluorophores is located at an internal site and the other fluorophore is attached to a terminus of the polynucleotide.

Dyes employed as quenchers include fluorescent dyes whose fluorescent characteristics are substantially unaffected by the presence or association with nucleic acids, particularly double stranded DNA. A reporter-quencher pair for a particular probe is selected such that the emission spectrum of the reporter overlaps with the absorption of the quencher (Figure 5). Spectral overlap allows for efficient energy transfer (FRET) when the probe is intact. Such dyes may include virtually any fluorescent dye fulfilling this criterion which is also

spectrally resolvable from whatever fluorophores that are employed on reporter-quencher probes. Dyes suitable as reporters may also be suitable as quenchers. Similarly, dyes suitable as quenchers may also be suitable as reporters. In one embodiment of a self-quenching probe, 6-carboxy-fluorescein (6-FAM) is used
5 as the reporter and 6-carboxytetramethylrhodamine (TAMRA) is used as the quencher such that the TAMRA dye substantially quenches any fluorescent emissions by 6-FAM.


In addition to DNA and RNA oligonucleotides, various nucleic acid analogs may be employed as self-quenching probes, such as, (i) oxygens in the
10 internucleotide linkage may be substituted by sulfur, carbon, or nitrogen, and (ii) the internucleotide phosphodiester linkages may be sulfate, carboxylate, amide, and the like. Alternatively, probes may have sugar modifications in one or more of the nucleosides, such as 2'-O- alkyl ribonucleosides and the like. Also, probes may have nucleobase modifications in one or more of the nucleosides, such as C-
15 5-alkynyl pyrimidines and the like. The fluorophores may be joined to the oligonucleotide or nucleic acid analog by appropriate functionalization of the fluorophores and/or the polymer building blocks. Detailed description of how to join fluorophores to nucleic acids and analogs can be found liberally in the literature (Andrus, 1992; Andrus, 1995; Hermanson, 1996; Ju et al, 1995).

20 The reporters and quenchers of self-quenching probes can be covalently attached to predetermined nucleotides of an oligonucleotide by using nucleoside phosphoramidite monomers containing reactive groups. For example, such reactive groups can be on a phosphate, or phosphate analog (Agrawal, S. 1990), on the 5'

hydroxyl when attachment is to the 5' terminal nucleotide (Andrus, 1995), and on base moieties (e.g. Ruth; Urdea et al; or the like). In further preference, the 3' terminal nucleotide of the oligonucleotide probe is blocked or rendered incapable of extension by a nucleic acid polymerase. Such 3' blocking is conveniently carried out by chemical attachment of a phosphate group (Horn, 1986; commercially available as PhosphaLink, PE Applied Biosystems). In a preferred embodiment of the invention, TAMRA is used as the quencher on self-quenching probes and the TAMRA dye is attached at the 3' terminus (Mullah, 1997; Mullah, 1998).

b. Block

10 The preferred block is a peptide-nucleic acid oligomer (PNA), a DNA analog in which the natural phosphodiester-deoxyribose backbone has been replaced by N-(2-aminoethyl)-glycine, a peptide-like unit (Nielsen, 1991). The nucleobases from natural nucleic acids, DNA and RNA, are retained through an amide linkage to the neutral backbone. The structure of PNA is represented below, 15 where B₁ and B₂ are nucleobases.



PNA oligomers are capable of recognizing complementary sequences by Watson/Crick base-pairing. Binding of PNA to its complement can occur in either a parallel or anti-parallel orientation of PNA; however, the anti-parallel

duplex is much more stable (Egholm et al). The melting temperature of PNA/DNA and PNA/RNA hybrids are much higher than corresponding DNA/DNA or DNA/RNA duplexes due to a lack of electrostatic repulsion in the PNA containing duplexes. PNA oligomers can be synthesized by conventional methods on commercially available, automated synthesizers, e.g. the Model 394 (PE Applied Biosystems, Foster City, CA), with commercially available reagents (Vinayak, 1997; Van der Laan, 1997).

IV. KIT OF REAGENTS

An aliquot from a master mix of a kit comprised of an internal control polynucleotide, internal control primers, a non-extending oligonucleotide or nucleic acid analog complementary to the internal control polynucleotide, target primers, a polymerase having 5'→3' nuclease activity, self-quenching fluorescence probes, 5' nucleotide triphosphates, and other reagents necessary for the exonuclease assay is delivered to all sample wells.

Embodiments of the invention include reagent compositions for use in nucleic acid amplification reactions. The subject compositions comprise a nucleic acid amplification buffer. The term "nucleic acid amplification buffer" as used herein, refers to a buffered aqueous solution that supports the enzymatic reaction or reactions required for a nucleic acid amplification reaction. The choice of buffer composition will vary in accordance with the particular enzyme selected for catalyzing the nucleic acid amplification reaction of interest (McPherson et al, 1991; McPherson et al, 1995; Dieffenbach, C., 1995). An example of a suitable nucleic

acid amplification buffer for Taq DNA polymerase catalyzed amplification reactions is: 10 mM Tris (pH 8.4), 50 mM KCl, 1.5 mM MgCl₂, 0.01% gelatin, 0.01% NP40, and 0.01% Tween™.

The reagent compositions of the invention may be supplied in a concentrated form or in a form that does not require significant dilution prior to use. The reagent compositions may be used by adding additional compounds required for performing the assay of interest, such compounds include, a thermostable polymerase, nucleotides, target polynucleotide, a reporter-quencher probes, and the like. After the addition of the necessary additional compounds, the reaction mixture may then be processed accordingly, e.g., thermocycling, so as to produce the desired amplification results.

V. DETECTION OF PCR PRODUCTS

The real-time detection system includes optical components operationally associated with a closed reaction chamber, which comprises a lens for focusing an excitation beam into the reaction mixture and for collecting the resulting fluorescence and a fiber optic for transmitting both the excitation beam from a light source to the lens and the fluorescent signals from the lens to a detection and analysis means. To induce fluorescence during amplification, laser light can be distributed to 96 sample wells via a multiplexed array of optical fibers. The resulting fluorescent emission returns via the fibers and is directed to a spectrograph with a charge-coupled device (CCD) camera. Preferably, the reaction mixture is contained in a closed reaction chamber to prevent

cross-sample contamination, or "carryover." The lens therefore focuses the excitation beam and collects fluorescence through a portion of a wall of the closed reaction chamber. The preferred reaction chamber is a tube, e.g. having the geometry and volume of a conventional microcentrifuge tube. The tube is closed after the reaction mixture is added by attaching a cap to the open end of the tube such that a leak-proof seal is formed. In a preferred embodiment of the sample interface for PCR, the lens directs the excitation beam and collects fluorescence generated by the probes through the cap of the tube. In other words, once a reaction mixture is placed in the tube and the cap is attached a closed reaction chamber is formed. Potential variability that could arise from sequential analysis of the first and second fluorescent signals is eliminated by simultaneously analyzing the signals by spectrally separating the signal light onto an array of photo detectors, e.g. by diffracting the signal onto a CCD array. An excitation beam generated by a single light source, e.g. a laser, is conveniently distributed to a plurality of closed reaction chambers by fiber optics. Likewise, the same fiber optics can collect the fluorescent signals from the plurality of reaction chambers for analysis by a single detection and analysis system. Alternatively the reaction chamber can be a well, e.g., a microtitre well, or a depression on a solid array. The array may be comprised of a plurality of spatially-addressable locations. Preferably, the system is employed with the PCR amplification of nucleic acids. An example of the preferred embodiment is the ABI PRISM™ 7700 Sequence Detection System (PE Applied Biosystems). Another embodiment is the ABI PRISM™ 7200 Sequence Detection System for

end-point analysis. Preferably, the system of the invention is employed to monitor PCR, although it may also be employed with other amplification schemes.

VI. DATA ANALYSIS

5 The results of an exonuclease assay can be analyzed using two parameters; the Rn value and the Ct value. The Rn value is the ratio of the fluorescence of a reporter dye and the fluorescence of the passive reference at the end of a PCR experiment. The Rn is calculated for each of the target and the ICP:

10 $Rn(\text{target}) = \text{Emission intensity of target probe} \div \text{Emission intensity of passive reference}$

$Rn(\text{ICP}) = \text{Emission intensity of ICP probe} \div \text{Emission intensity of passive reference}$

15 The Ct value, or threshold cycle number, is the PCR cycle number at which the fluorescence ratio is distinguishable from the background. For a given reporter dye and a fixed concentration of target, both the Rn and Ct values reflect the efficiency of the quencher.

VII. EXAMPLES

20 The invention will be further clarified by a consideration of the following examples, which are intended to be purely exemplary of the present invention and not to in any way limit its scope.

EXAMPLE 1**Preparation of doubly-labeled, self-quenching probe for exonuclease assay**

Automated synthesis of self-quenching probes can be performed using an Applied Biosystems Model 394 DNA/RNA synthesizer (The Perkin-Elmer Corporation, PE Applied Biosystems Division) according to the general procedures described in the users manual. A 5' FAM, 3' TAMRA probe can be synthesized at 0.2 μ mol scale using TAMRA-labeled CPG solid supports (Mullah et al, 1997; Mullah et al, 1998), the set of phosphoramidite nucleosides A^{bz}, G^{dmf}, C^{bz}, T, other reagents recommended by the manufacturer (PE Applied Biosystems) and FAM dye-labeled phosphoramidite (Theisen et al). The standard 0.2 μ mol synthesis cycle is slightly modified by extending coupling time of FAM amidite by an additional 120 seconds. Each probe includes a reporter dye attached to a 5'-end of and a quencher dye located at a 3'-end of the probe (Figure 6).

After completion of the synthesis, oligonucleotides are autocleaved from the support on the DNA synthesizer by treating with a mixture of MeOH:t-BuNH₂:H₂O (1:1:2) (Woo, 1993) for a 1 hr autocleavage procedure as described in the operators manual for the Applied Biosystems Model 394 DNA/RNA synthesizer. Base protecting groups are removed by heating the mixture at 85°C for 1 hr or at 65°C for 3 h. The oligonucleotides can be analyzed and purified by reverse phase HPLC, anion-exchange HPLC, capillary gel electrophoresis, polyacrylamide gel electrophoresis, and other conventional techniques (Andrus, 1992).

EXAMPLE 2

Detection of C-myc mRNA by Exonuclease Assay using Exogenous IPC,

Reverse-transcription

5 The cDNA is prepared using 50 ng human RNA and C-MYC reverse primer:

5' CAACATCGAT TTCTTCCTCA TCTTCT 3' (SEQ. ID NO. 1)

in 100 µl solution containing 1X PCR buffer II, 5.5 mM MgCl₂, 500 µM each of dATP, dCTP, dTTP, and dGTP, 1.25 U/µl of MuLV reverse transcriptase, 0.4
10 U/µl RNase inhibitor, and RNase free H₂O at 48°C for 30 min in an ABI Model 9600 thermal cycler.

Reagents

1) The following kit of reagents is mixed and 45 µl is dispensed into each of all 96 wells, A1-H12 (Figure 7):

15 Tris-HCl pH 8.0, 100 mM, 16% glycerol, 0.1% gelatin, 0.02%, Tween 20™, 10 mM MgCl₂,
400 µM each of dATP, dCTP, deaza dGTP and 800 µM of dUTP,
0.1U/ µl of AmpliTaqGold DNA polymerase,
0.02 U/ µl of Amperase UNG,
20 120 nM of Passive Reference,
forward C-MYC target primer:

5' GCCCCTGGTT GCTCCATGA 3' (SEQ. ID NO. 2)

reverse C-MYC target primer

100 nM FAM labeled self-quenching target probe:

5' FAM- CAGCACAACT ACGCAGCGCC TCCT -TAMRA 3'

(SEQ. ID NO. 3)

Exogenous Internal Positive Control (IPC) reagents:

5 Plasmid DNA harboring a 80 bp amplicon internal control
polynucleotide (IPC)

50 nM forward IPC primer:

5' TCCAACCGCC AACTATCAA 3' (SEQ. ID NO. 4)

50 nM reverse IPC primer:

10 5' CATCCGCACA CTATCTCATC GT 3' (SEQ. ID NO. 5)

200 nM of self-quenching IPC probe:

5' JOE- CAGCTCGTTG ATCTCCGTT CTGGCA -

TAMRA 3' (SEQ. ID NO. 6)

2) Ten µl of the cDNA is added to each of 84 wells, B1-H12 (Figure 7)

15 3) PNA block is added to wells A1-A6 (Figure 7):

300 nM PNA block:

H₂N - TCCAACCCGC CACCCACTAT C - CONH₂ (SEQ.

ID NO. 7)

PCR

20 Human cDNA is amplified by thermal cycling conditions that begin with
2 min at 50°C, 10 min hold at 95°C and then 40 cycles of: 15 sec denaturation at
95°C and 1 min annealing and extension at 60°C. Thermal cycling and real-time

fluorescence detection is conducted on an ABI PRISM™ 7700 Sequence Detection System (Perkin-Elmer Co.)

The target cDNA and IPC are amplified in the presence of the IPC, detected by the presence of FAM and JOE signals respectively in wells B1-H12.

- 5 The PNA block inhibits the amplification of the IPC in wells A1-A6, detected by the absence of JOE signal. No target amplification is detected, by the lack of FAM signal, in wells A1-A12. IPC is amplified, as detected by JOE signal, in wells A7-A12 (Figure 7).

EXAMPLE 3

- 10 Detection of DNA: NPTII gene from transgenic cotton plants by exonuclease assay with exogenous IPC

Approximately 200 ng transgenic cotton genomic DNA is prepared by standard techniques for the 96 well test,

Reagents

- 15 The same kit of reagents as Example 2, including the same IPC reagents, is mixed and 45 µl total volume is dispensed into each of all 96 wells, A1-H12 (Figure 7), including NPTII target specific primers and probe:

300 nM forward target primer:

5' CAGGACGGGC GTTCCTTGC 3' (SEQ. ID NO. 8)

- 20 300 nM reverse target primer:

5' GTGGGTCGAA TGGGCAGGTAG C 3' (SEQ. ID NO. 9)

200 nM self-quenching target probe:

5' FAM - ACTGAAGCGG GAAGGGACTG GCT - TAMRA 3'

(SEQ. ID NO. 10)

PCR

5 Amplification is conducted under the same conditions and on the same system as Example 2.

Both the cotton sample DNA and the IPC are successfully amplified. Comparison of the Ct with and without the IPC suggest that PCR efficiency is not affected by the complexity of the sample DNA.

10

EXAMPLE 4

Detection of E. Coli 0157:H7 by exonuclease assay with exogenous IPC

Approximately 100 ng. DNA from purified plasmid with cloned target sequence
15 is prepared by standard techniques for the 96 well test. A range 10,000 to less than 1 copies per well are used in the unknown samples.

Reagents

The same kit of reagents as Example 2, including the same IPC reagents, is mixed and 45 µl total volume is dispensed into each of all 96 wells, A1-H12
20 (Figure 7), including NPTII target specific primers and probe:

300 nM forward target primer:

5' ATCCTGGACC AGGTTCTCTGA 3' (SEQ. ID NO. 11)

300 nM forward target primer:

5' CGGTACAAGC TGCAACTGTTA 3' (SEQ. ID NO. 12)

200 nM Target probe:

5' FAM- TAGGCAGTAT TCCGAAATGAC - TAMRA 3' (SEQ. ID NO.

5 13)

PCR

Amplification is conducted under the same conditions and on the same system as Example 2.

The results are shown in Figure 8. The unknown samples of 1 to 10,000
10 copies gave positive detection by detection of the FAM signal above background.

EXAMPLE 5

Detection of Mycoplasma synoviae DNA extracted from cultured cells by
15 exonuclease assay with exogenous IPC

Approximately 100 ng. DNA from multiple sources of Mycoplasma synoviae is prepared by standard techniques for the 96 well test. Each source is tested as ten-fold replicates.

Reagents

20 The same kit of reagents as Example 2, including the same IPC reagents is mixed and 45 µl total volume is dispensed into each of all 96 wells, A1-H12 (Figure 7).

The kit of reagents includes:

Forward Primer:

5' CGGATTGTAG TCTGCAACTCGACT A 3' (SEQ. ID NO. 14)

Reverse Primer:

5' ACAAACCGAC TTCGGGCATT 3' (SEQ. ID NO. 15)

5 Self-quenching probe:

5' FAM - AATACGTTCT CGGGTCTTGT ACACACCGC - TAMRA

3' (SEQ. ID NO. 16)

PCR

Amplification is conducted under the same conditions and on the same
10 system as Example 2. The target amplicon length is 143 bp.

The results are shown in Figure 9. All No Amplification Control samples
(A1-A6), containing IPC, PNA block and no target, gave insignificant FAM or
JOE signals, as expected. All No Template Control samples (A7-A12),
containing IPC, no PNA block and no target, gave insignificant FAM and
15 significant JOE signal, as expected. All Unknown samples (B1-H12), containing
IPC, no PNA block, and target, gave true positive results, except for sample H8
(target Ct=40) indicating true negative.

20 All publications and patent applications are herein incorporated by
reference to the same extent as if each individual publication or patent
application was specifically and individually indicated to be incorporated by
reference.

Although only a few embodiments have been described in detail above, those having ordinary skill in the molecular biology art will clearly understand that many modifications are possible in the preferred embodiment
5 without departing from the teachings thereof. All such modifications are intended to be encompassed within the following claims.

WE CLAIM:

1. A method of conducting nucleic acid amplification reactions in a single reaction chamber comprising:

5 providing an internal control polynucleotide, internal control primers, a non-extendable oligonucleotide or oligonucleotide analog complementary to the internal control polynucleotide, a target polynucleotide, target primers, a nucleic acid polymerase having 5'→3' nuclease activity, 5' nucleotide triphosphates; self-quenching fluorescence probes including reporter and quencher moieties,

10 said probes existing in at least one single-stranded conformation when unhybridized wherein said quencher dye quenches the fluorescence of said reporter dye and at least one conformation when hybridized to said polynucleotides, wherein the fluorescence of the reporter dye is unquenched, the fluorescence intensity of reporter dye being greater than
15 the fluorescence intensity of quencher dye when probe is hybridized to polynucleotide; and the steps of

hybridizing said target primers to said target polynucleotide;

hybridizing said internal control primers to said internal control polynucleotide;

20 amplifying the target polynucleotide by PCR, whereby target polynucleotide amplification products are produced; and

amplifying the internal control polynucleotide by PCR, whereby internal control polynucleotide amplification products are produced.

2. The method of claim 1 wherein said nucleic acid polymerase digests said probes during amplification to separate said reporter dye from said quencher dye.

3. The method of claim 1 wherein a first self-quenching fluorescence probe is complementary to the internal control polynucleotide and a second self-quenching fluorescence probe is complementary to the target polynucleotide.

4. The method of claim 3 wherein the reporter dye of the first self-quenching fluorescence probe is spectrally resolvable from the reporter dye of the second self-quenching fluorescence probe.

5. The method of claim 1 wherein a product of the nucleic acid amplification of the internal control polynucleotide is spectrally resolvable from the products of the nucleic acid amplification of the target polynucleotide.

6. The method of claim 1 wherein the products of the nucleic acid amplification of the internal control polynucleotide and target polynucleotide are measured and quantitated by end-point analysis.

7. The method of claim 1 wherein the products of nucleic acid amplification of the internal control polynucleotide and target polynucleotide are measured and quantitated by real-time analysis.





8. The method of claim 1 wherein the products of nucleic acid amplification of the internal control polynucleotide and target polynucleotide are measured by fluorescence detection.

9. The method of claim 1 wherein said nucleic acid polymerase is a
5 thermal-stable polymerase with exonuclease activity.

10. The method of claim 1 wherein said reporter is a xanthene dye.


11. The method of claim 10 wherein said xanthene dye is a fluorescein dye.

12. The method of claim 11 wherein said fluorecein dye is selected from
- 10 the group consisting of



where L is a linker, and including substituted forms thereof.

13. The method of claim 1 wherein said quencher is selected from the group consisting of




5 where L is a linker, and including substituted forms thereof.

14. The method of claim 1 wherein said reporter dye is separated from said quencher dye by at least 12 nucleotides.


15. The method of claim 1 wherein said reporter dye is attached at the 5' terminus or 3' terminus of the self-quenching fluorescence probe.

10 16. The method of claim 1 wherein said quencher dye is attached at the 5' terminus or 3' terminus of the self-quenching fluorescence probe.

17. The method of claim 1 wherein said non-extending oligonucleotide or nucleic acid analog complementary to the internal control polynucleotide is selected from the group consisting of



where R is fluoro, chloro, amino, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}=\text{CH}_2$, and $-\text{OCH}_2\text{CH}_2\text{OCH}_3$,



including substituted forms thereof.

18. A kit of reagents for nucleic acid amplification comprising:

an internal control polynucleotide, internal control primers, a non-extending oligonucleotide or nucleic acid analog complementary to the internal control polynucleotide, target primers, a nucleic acid polymerase having 5'→3' nuclease activity, self-quenching fluorescence probes including reporter and quencher moieties, 5' nucleotide triphosphates; and other reagents necessary for nucleic acid amplification.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: The Perkin-Elmer Corporation
- (ii) TITLE OF THE INVENTION: METHODS FOR EXOGENOUS, INTERNAL CONTROLS DURING NUCLEIC ACID AMPLIFICATION
- (iii) NUMBER OF SEQUENCES: 16
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: The Perkin-Elmer Corporation
 - (B) STREET: 850 Lincoln Centre Drive
 - (C) CITY: Foster City,
 - (D) STATE: California
 - (E) COUNTRY: USA
 - (F) ZIP: 94404
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: To Be Assigned
 - (B) FILING DATE: 26-MAR-1998
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 09/048,880
 - (B) FILING DATE: March 26, 1998
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Scott R. Bortner
 - (B) REGISTRATION NUMBER: 34,298
 - (C) REFERENCE/DOCKET NUMBER: 4382WO
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (650) 638-6245
 - (B) TELEFAX: (650) 638-6071

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAACATCGAT TTCTTCCTCA TCTTCT

26

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCCCCCTGGTT GCTCCATGA

19

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CAGCACAACT ACGCAGCGCC TCCT

24

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TCCAACCGCC ACACTATCAA

20

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TCCAACCGCC ACACTATCAA

20

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CAGCTCGTTG ATCTTCCGTT CTGGCA

26

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TCCAACCCGC CACCCACTAT C

21

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CAGGACGGGC GTTCCTTGC

19

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GTGGGTCGAA TGGGCAGGTA GC

22

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACTGAAGCGG GAAGGGACTG GCT

23

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATCCTGGACC AGGTTCTGA

20

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CGGTACAAGC TGCAACTGTT A

21

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TAGGCAGTAT TCCGAAATGA C

21

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGGATTGTAG TCTGCAACTC GACTA

25

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ACAAACCGAC TTCGGGCATT

20

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AATACGTTCT CGGGTCTTGT ACACACCGC

29




Figure 1







Figure 2

3/11







Figure 3

4/11




Figure 4

5/11

Dye	Emission maxima (nm)	Absorbance maxima (nm)
5-FAM	522	494
6-FAM	518	492
JOE	554	528
HEX	553	535
TET	538	521
TAMRA	582	560
ROX	607	587

Figure 5.

6/11




Figure 6

7/11

96 Well, microtitre plate:

	1	2	3	4	5	6	7	8	9	10	11	12
A	IPC-	IPC-	IPC-	IPC-	IPC-	IPC-	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
B	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
C	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
D	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
E	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
F	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
G	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+
H	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+	IPC+

Sample assignment:

wells	reagents	function	sample signal	IPC signal
A1-A6	IPC, block	no amplification control	-	-
A7-A12	IPC	no template control	-	+
B1-H12	targets, IPC	unknowns, samples	+ or -	+

Figure 7

8/11

Well	Type	Sample	PCR	target ΔR_n	target C_t
A1	NAC		No amp	0.06	40
A2	NAC		No amp	0.06	40
A3	NAC		No amp	0.06	40
A4	NAC		No amp	0.06	40
A5	NAC		No amp	0.06	40
A6	NAC		No amp	0.06	40
A7	NTC		-	0.05	40
A8	NTC		-	0.05	40
A9	NTC		-	0.06	40
A10	NTC		-	0.04	40
A11	NTC		-	0.06	40
A12	NTC		-	0.05	40
B1	Unknown	10,000	+	1.72	26.38
B2	Unknown	1000	+	1.45	29.57
B3	Unknown	100	+	0.81	33.04
B4	Unknown	10	+	0.12	38.01
B5	Unknown	1	+	0.14	37.94
B6	Unknown	0.1	-	0.05	40
B7	Unknown	Negative	-	0.05	40
B8	Unknown	Negative	-	0.05	40
B9	Unknown	Negative	-	0.05	40
B10	Unknown	Negative	-	0.04	40
B11	Unknown	Negative	-	0.05	40
B12	Unknown	Negative	-	0.05	40
C1	Unknown	Negative	-	0.05	40
C2	Unknown	Negative	-	0.04	40
C3	Unknown	Negative	-	0.04	40
C4	Unknown	Negative	-	0.03	40
C5	Unknown	Negative	-	0.04	40
C6	Unknown	Negative	-	0.04	40
C7	Unknown	Negative	-	0.04	40
C8	Unknown	Negative	-	0.04	40
C9	Unknown	Negative	-	0.04	40
C10	Unknown	Negative	-	0.04	40
C11	Unknown	Negative	-	0.05	40
C12	Unknown	Negative	-	0.04	40
D1	Unknown	10,000	+	1.66	26.25
D2	Unknown	1000	+	1.35	30.17
D3	Unknown	100	+	0.72	33.09
D4	Unknown	10	+	0.28	36.09
D5	Unknown	1	-	0.05	40
D6	Unknown	0.1	-	0.04	40
D7	Unknown	Negative	-	0.04	40
D8	Unknown	Negative	-	0.04	40
D9	Unknown	Negative	-	0.04	40
D10	Unknown	Negative	-	0.04	40
D11	Unknown	Negative	-	0.05	40
D12	Unknown	Negative	-	0.05	40
E1	Unknown	Negative	-	0.05	40
E2	Unknown	Negative	-	0.04	40
E3	Unknown	Negative	-	0.04	40
E4	Unknown	Negative	-	0.05	40

Figure 8

RECTIFIED SHEET (RULE 91)

ISA/EP

9/11

E5	Unknown	Negative	-	0.04	40
E6	Unknown	Negative	-	0.05	40
E7	Unknown	Negative	-	0.05	40
E8	Unknown	Negative	-	0.05	40
E9	Unknown	Negative	-	0.05	40
E10	Unknown	Negative	-	0.05	40
E11	Unknown	Negative	-	0.05	40
E12	Unknown	negative	-	0.05	40
F1	Unknown	10,000	+	1.65	40
F2	Unknown	1000	+	1.37	40
F3	Unknown	100	+	0.72	40
F4	Unknown	10	+	0.67	40
F5	Unknown	1	-	0.05	40
F6	Unknown	0.1	-	0.04	40
F7	Unknown	negative	-	0.05	40
F8	Unknown	negative	-	0.05	40
F9	Unknown	negative	-	0.05	40
F10	Unknown	negative	-	0.05	40
F11	Unknown	negative	-	0.05	40
F12	Unknown	negative	-	0.04	40
G1	Unknown	negative	-	0.05	40
G2	Unknown	negative	-	0.04	40
G3	Unknown	negative	-	0.06	40
G4	Unknown	negative	-	0.05	40
G5	Unknown	negative	-	0.04	40
G6	Unknown	negative	-	0.04	40
G7	Unknown	negative	-	0.05	40
G8	Unknown	negative	-	0.04	40
G9	Unknown	negative	-	0.05	40
G10	Unknown	negative	-	0.05	40
G11	Unknown	negative	-	0.05	40
G12	Unknown	negative	-	0.04	40
H1	Unknown	negative	-	0.05	40
H2	Unknown	negative	-	0.04	40
H3	Unknown	negative	-	0.05	40
H4	Unknown	negative	-	0.04	40
H5	Unknown	negative	-	0.04	40
H6	Unknown	negative	-	0.04	40
H7	Unknown	negative	-	0.04	40
H8	Unknown	negative	-	0.05	40
H9	Unknown	negative	-	0.04	40
H10	Unknown	negative	-	0.04	40
H11	Unknown	negative	-	0.04	40
H12	Unknown	negative	-	0.04	40

Figure 8 (Continued)

10/11

Well	Type	target ΔR_n	Target Ct	ICP ΔR_n	ICP Ct
A1	NAC	0.005	40	0.009	40
A2	NAC	0.0007	40	0.01	40
A3	NAC	0.01	40	0.001	40
A4	NAC	0.002	40	0.01	40
A5	NAC	0.003	40	0.002	40
A6	NAC	0.005	40	0.0005	40
A7	NTC	0.02	31.88	0.51	24.89
A8	NTC	0.04	32.71	0.52	25.02
A9	NTC	0.02	35.72	0.52	24.41
A10	NTC	0.03	30.96	0.53	25.02
A11	NTC	0.03	30.47	0.52	24.61
A12	NTC	0.04	31.12	0.52	25.04
B1	Unknown	1.41	17.1	0.54	24.35
B2	Unknown	1.43	17.13	0.56	23.15
B3	Unknown	1.43	16.93	0.52	25.65
B4	Unknown	1.36	16.87	0.52	24.05
B5	Unknown	1.40	17.18	0.51	25.45
B6	Unknown	1.41	16.93	0.50	23.75
B7	Unknown	1.43	16.87	0.52	26.41
B8	Unknown	1.43	17.18	0.51	23.47
B9	Unknown	1.36	16.83	0.55	25.03
B10	Unknown	1.40	16.95	0.51	23.28
B11	Unknown	1.41	17.20	0.54	25.91
B12	Unknown	1.46	16.97	0.50	22.65
C1	Unknown	1.41	16.73	0.53	25.03
C2	Unknown	1.41	17.02	0.54	24.64
C3	Unknown	1.39	20.79	0.51	24.52
C4	Unknown	1.44	20.91	0.52	23.40
C5	Unknown	1.35	21.17	0.54	24.98
C6	Unknown	1.26	21.02	0.54	24.35
C7	Unknown	1.25	20.88	0.53	24.92
C8	Unknown	1.26	21.15	0.53	24.88
C9	Unknown	1.20	21.00	0.52	25.01
C10	Unknown	1.28	21.19	0.51	24.72
C11	Unknown	1.24	21.01	0.51	24.64
C12	Unknown	1.25	21.05	0.50	25.01
D1	Unknown	1.26	21.07	0.53	24.58
D2	Unknown	1.20	20.79	0.51	25.16
D3	Unknown	1.26	21.05	0.51	24.97
D4	Unknown	1.20	21.07	0.49	24.94
D5	Unknown	0.98	25.33	0.53	24.87
D6	Unknown	1.22	21.32	0.51	24.99
D7	Unknown	1.04	25.05	0.51	25.04
D8	Unknown	1.03	24.33	0.48	25.25
D9	Unknown	1.00	25.21	0.49	25.27
D10	Unknown	1.01	25.29	0.51	25.36
D11	Unknown	0.99	25.20	0.53	24.58
D12	Unknown	1.03	24.31	0.50	24.97
E1	Unknown	1.00	25.17	0.51	25.35
E2	Unknown	1.04	24.98	0.49	24.58
E3	Unknown	1.04	24.86	0.49	24.97
E4	Unknown	1.01	24.64	0.40	25.35

Figure 9

RECTIFIED SHEET (RULE 91)
ISA/EP

11/11

E5	Unknown	0.73	28.31	0.50	25.28
E6	Unknown	0.74	28.51	0.50	25.81
E7	Unknown	0.76	28.31	0.49	25.56
E8	Unknown	0.76	28.51	0.49	25.54
E9	Unknown	0.60	28.49	0.50	25.55
E10	Unknown	0.74	28.45	0.50	26.01
E11	Unknown	0.72	28.25	0.48	25.54
E12	Unknown	0.75	29.48	0.49	26.02
F1	Unknown	0.77	30.11	0.48	25.30
F2	Unknown	0.38	30.52	0.48	25.30
F3	Unknown	0.41	29.87	0.51	25.16
F4	Unknown	0.37	33.83	0.47	25.43
F5	Unknown	0.39	31.79	0.48	25.77
F6	Unknown	0.40	32.06	0.48	25.90
F7	Unknown	0.41	31.16	0.50	26.03
F8	Unknown	0.39	31.70	0.50	25.46
F9	Unknown	0.41	31.29	0.48	25.53
F10	Unknown	0.39	32.20	0.49	25.83
F11	Unknown	0.41	31.73	0.50	25.91
F12	Unknown	0.41	33.61	0.50	25.32
G1	Unknown	0.39	30.86	0.48	25.32
G2	Unknown	0.41	35.58	0.49	25.44
G3	Unknown	0.39	32.36	0.50	25.57
G4	Unknown	0.41	33.43	0.50	25.51
G5	Unknown	0.40	29.50	0.48	25.55
G6	Unknown	0.13	33.55	0.49	25.35
G7	Unknown	0.086	29.86	0.48	25.28
G8	Unknown	0.12	31.29	0.49	25.53
G9	Unknown	0.088	32.20	0.49	25.01
G10	Unknown	0.12	31.73	0.50	25.38
G11	Unknown	0.10	33.61	0.52	25.45
G12	Unknown	0.14	30.86	0.51	25.52
H1	Unknown	0.05	35.58	0.51	25.04
H2	Unknown	0.03	32.36	0.51	24.87
H3	Unknown	0.03	33.43	0.52	24.65
H4	Unknown	0.04	29.50	0.52	25.31
H5	Unknown	0.045	33.55	0.51	24.84
H6	Unknown	0.055	29.86	0.51	25.14
H7	Unknown	0.040	29.57	0.49	25.15
H8	Unknown	0.0009	40	0.51	24.97
H9	Unknown	0.042	30.94	0.52	25.20
H10	Unknown	0.05	30.29	0.48	25.17
H11	Unknown	0.05	30.18	0.54	24.88
H12	Unknown	0.04	31.26	0.54	25.19

Figure 9 (Continued)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number
WO 01/16367 A1

- (51) International Patent Classification⁶: C12Q 1/68 (81) Designated States (*national*): AU, CA, JP.
- (21) International Application Number: PCT/US99/06978 (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (22) International Filing Date: 27 March 1999 (27.03.1999)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant: THE PERKIN-ELMER CORPORATION [US/US]; 850 Lincoln Centre Drive, Foster City, CA 94404 (US).
- (48) Date of publication of this corrected version: 1 November 2001
- (72) Inventors: AOYAGI, Kazuko; 3 Admiral Drive #457, Emeryville, CA 94608 (US). LIVAK, Kenneth, J.; 1148 Polk Lane, San Jose, CA 95117 (US).
- (15) Information about Correction: see PCT Gazette No. 44/2001 of 1 November 2001, Section II
- (74) Agent: BORTNER, Scott, R.; The Perkin-Elmer Corporation, 850 Lincoln Centre Drive, Foster City, CA 94404 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR EXOGENOUS, INTERNAL CONTROLS DURING NUCLEIC ACID AMPLIFICATION



(57) Abstract: Reporter-quencher probe assays of nucleic acid amplification, such as PCR, are rendered more meaningful by the addition of internal control reagents. An internal control polynucleotide is amplified with internal control primers and the product is measured by correlation with increased fluorescence by polymerase mediated-exonuclease cleavage or hybridization of the internal control probe. Probes specific for target and internal control polynucleotides are labelled with spectrally resolvable reporters, allowing for concurrent detection and measurement of target and control amplification. A kit of all PCR reagents can be dispensed into reaction chambers in a high-throughput system for rapid and accurate nucleic acid amplification assay, with real-time or end-point measurements. Fluorescent signals correlated to target and internal control levels are spectrally resolvable and measured concurrently. A non-extending oligonucleotide or nucleic analog "block", complementary to the internal control polynucleotide, is added to the amplification mixture to preclude amplification of the internal control polynucleotide and function as an internal negative control. The amplification control reagents, kits, and methods of the present invention provide positive

and negative control tests occurring within, and measurable within, the reaction chamber.

WO 01/16367 A1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06978

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GIBSON U E M ET AL: "A NOVEL METHOD FOR REAL TIME QUANTITATIVE RT-PCR" GENOME RESEARCH, vol. 6, no. 10, 1 October 1996 (1996-10-01), pages 995-1001, XP000642796 ISSN: 1088-9051 the whole document	1-18
Y	HEID C A ET AL: "REAL TIME QUANTITATIVE PCR" GENOME RESEARCH, vol. 6, no. 10, 1 October 1996 (1996-10-01), pages 986-994, XP000642795 ISSN: 1088-9051 the whole document	1-18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

22 July 1999

Date of mailing of the international search report

04/08/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Hagenmaier, S

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/06978

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	REISCHL U ET AL: "QUANTITATIVE PCR A SURVEY OF THE PRESENT TECHNOLOGY" MOLECULAR BIOTECHNOLOGY, vol. 3, 1 January 1995 (1995-01-01), pages 55-71, XP000600241 ISSN: 1073-6085 the whole document	1-18
Y	US 5 571 673 A (PICONE TERESA K H) 5 November 1996 (1996-11-05) See column 10, line 34- line 44. the whole document	1-18
Y	WO 92 02638 A (CETUS CORP) 20 February 1992 (1992-02-20) the whole document	1-18
Y	US 5 219 727 A (DOYLE MICHAEL V ET AL) 15 June 1993 (1993-06-15) the whole document	1-18
Y	WO 97 46708 A (PERKIN ELMER CORP) 11 December 1997 (1997-12-11) the whole document	1-18
Y	WOUDENBERG T M ET AL: "QUANTITATIVE PCR BY REAL TIME DETECTION" PROCEEDINGS OF THE SPIE, vol. 2680, 1 January 1996 (1996-01-01), pages 306-315, XP000197422 the whole document	1-18
Y	EP 0 672 677 A (HOECHST AG) 20 September 1995 (1995-09-20) the whole document	17
Y	VAN DER LAAN ET AL.: "A CONVENIENT AUTOMATED SOLID-PHASE SYNTHESIS OF PNA-(5')-DNA-(3')-PNA CHIMERA" TETRAHEDRON LETTERS, vol. 38, no. 13, 1997, pages 2249-2252, XP002110033 cited in the application the whole document	17
Y	US 5 516 663 A (BACKMAN KEITH C ET AL) 14 May 1996 (1996-05-14) column 3, line 53 - line 64	17

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/US 99/06978

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5571673 A	05-11-1996	AU 700885 B AU 3901095 A CA 2163388 A CN 1133888 A EP 0713921 A JP 8220092 A NZ 280506 A	14-01-1999 30-05-1996 24-05-1996 23-10-1996 29-05-1996 30-08-1996 27-07-1997
WO 9202638 A	20-02-1992	US 5210015 A AU 666837 B AU 8651091 A CA 2088683 A EP 0543942 A EP 0919565 A JP 2825976 B JP 6500021 T US 5804375 A US 5487972 A	11-05-1993 29-02-1996 02-03-1992 07-02-1992 02-06-1993 02-06-1999 18-11-1998 06-01-1994 08-09-1998 30-01-1996
US 5219727 A	15-06-1993	AT 178656 T AU 653920 B AU 6352390 A CA 2064906 A DE 69033044 D EP 0497784 A JP 5504886 T US 5476774 A WO 9102817 A	15-04-1999 20-10-1994 03-04-1991 22-02-1991 12-05-1999 12-08-1992 29-07-1993 19-12-1995 07-03-1991
WO 9746708 A	11-12-1997	US 5736333 A AU 3230297 A CA 2256457 A EP 0904411 A	07-04-1998 05-01-1998 11-12-1997 31-03-1999
EP 0672677 A	20-09-1995	DE 4408528 A AU 698210 B AU 1479895 A CA 2144475 A CN 1112126 A FI 951132 A JP 7278179 A NO 950955 A	28-09-1995 29-10-1998 21-09-1995 15-09-1995 22-11-1995 15-09-1995 24-10-1995 15-09-1995
US 5516663 A	14-05-1996	US 5427930 A AU 6634294 A CA 2160784 A EP 0695367 A JP 8509125 T WO 9424311 A AT 137269 T AU 635105 B AU 7000791 A CA 2035010 A,C DE 69118930 D DE 69118930 T EP 0439182 A ES 2089038 T JP 1980321 C	27-06-1995 08-11-1994 27-10-1994 07-02-1996 01-10-1996 27-10-1994 15-05-1996 11-03-1993 01-08-1991 27-07-1991 30-05-1996 09-01-1997 31-07-1991 01-10-1996 17-10-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/06978

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5516663 A		JP 4211399 A	03-08-1992
		JP 6036760 B	18-05-1994
		KR 9513953 B	18-11-1995
		KR 9602235 B	13-02-1996
		US 5573907 A	12-11-1996
		US 5792607 A	11-08-1998
		CA 2111962 A	07-01-1993
		EP 0596918 A	18-05-1994
		JP 6508750 T	06-10-1994
		WO 9300447 A	07-01-1993
		US 5453355 A	26-09-1995
		AU 3943093 A	08-11-1993
		WO 9320191 A	14-10-1993
		AU 4687393 A	03-03-1994
		CA 2140331 A	17-02-1994
		EP 0654093 A	24-05-1995
		JP 8501212 T	13-02-1996
		WO 9403636 A	17-02-1995

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.